

<<大豆育种的分子标记系统最新进展>>

图书基本信息

书名：<<大豆育种的分子标记系统最新进展>>

13位ISBN编号：9787030346780

10位ISBN编号：7030346785

出版时间：2012-1

出版时间：科学出版社

作者：李文彬、David A.Lightfoot

页数：263

字数：567000

版权说明：本站所提供下载的PDF图书仅提供预览和简介，请支持正版图书。

更多资源请访问：<http://www.tushu007.com>

## <<大豆育种的分子标记系统最新进展>>

### 内容概要

现代分子标记技术与大豆育种（英文版）系统地介绍了各种类型分子标记的发展概况和开发技术，QTL分析和分子辅助育种的基本原理，阐述了大豆产量性状、品质性状、抗病虫性状、动态发育性状分子标记和QTL的最新研究进展及大豆分子设计育种的基本设想及展望。该书反映了近年分子标记在大豆遗传改良方面获得的成果。

作者简介

Wenbin Li

Key Laboratory of Soybean Biology in Chinese Ministry of Education(Agriculture Research System of China and the Key Laboratory of Northeastern Soybean Biology and Breeding/Genetics of Chinese Agriculture Ministry),Northeast Agricultural University,Harbin,China.150030

David A.Lightfoot

Illinois Soybean Center for Excellence in Soybean Research,Southern Illinois University at Carbondale,Carbondale,IL 62901,USA

书籍目录

Acknowledgement  
Preface  
Chapter 1 Essential principles of molecular marker and quantitative trait loci associations  
Chapter 2 Method development for identifying functional markers in the soybean genome  
Chapter 3 Map development and marker-assisted selection for oligogenic and polygenic traits  
Chapter 4 QTL analysis of fungal disease resistance  
Chapter 5 Identification of loci for resistance to insects and nematodes  
Chapter 6 QTL analysis for seed quality traits  
Chapter 7 Seed yield and component trait loci  
Chapter 8 Loci underlying soil edaphic, physiological and developmental traits  
Chapter 9 Dynamic QTL analysis of major quantitative traits  
Chapter 10 Molecular mapping and breeding for biotic stress resistance  
Colored Figures

## 章节摘录

A novel DNA-cleaving reagent became available when it was shown that Mu transposition preferentially targets mismatched sites in DNA ( Yanagihara and Mizuuchi 2002 ). This paved the way to establish the mismatch-targeting methodology and indicate, using a known polymorphic test fragment, that mutation indeed can be detected by the use of the Mu transposon approach. Mu A transposase protein assembles two transposon end segments into a tetrameric DNA transposition complex, which can capture the target DNA and cause the strand to transfer. The transposon DNA is joined into the target when the target DNA strands are simultaneously cleaved, where nearly 90% of the strand transfers occur when mismatches are present in the target DNA. DNA duplexes that contain mismatches can be obtained by amplification of regions where the genomic DNA contains at least two alleles, or when the region is amplified from a heterozygous individual or from a sample that combines DNA from two or more individuals representing different allelic variants ( Orsini et al. 2007 ).

Prospect for Future Study of Functional Markers Many candidate genes for functional marker development are available and allele sequencing projects are in process in soybean ( Afzal et al. 2012 ), Arabidopsis ( <http://walnut.usc.edu/> ) and other economic crop species ( e.g. <http://www.grasp-euv.dk/>; <http://www.maizegenetics.net/> ). However, even in model species, less than 10% of all genes have been functionally characterized, whereas this number is probably less than 5% in non-model species. Furthermore, the criteria by which such genes have been functionally characterized ( e.g. in a biological sense ) might not be sufficient to establish gene function in an agronomic sense. Thus, further functional characterization might be necessary before proceeding to functional marker development. Furthermore, a crucial question will be whether useful allelic variation can be identified for all genes of ecological and agronomical relevance. Given that functionally described target genes are available, the limiting step in functional marker development will be the choice and development of suitable plant materials and its thorough phenotypic characterization to distinguish minor phenotypic effects.

Refining the predictive value of functional markers will require the development of an experimental, statistical and bioinformatics framework for data accumulation and evaluation across different studies. The soybean genome data have been published ( Schmutz et al. 2010 ). The corresponding research base on the genome will rapidly develop, including functional markers ( Wu et al. 2010; Lam et al. 2010 ). Association studies and isogenic comparisons of homozygous genotypes can be extended to test crosses using different testers to determine both additive and dominance effects of functional sequence motifs. Estimates of additive and dominance effects are determined in a relative manner ( i.e. the estimates depend on the trait locus alleles as well as the genetic background of genotypes present in a specific study ). The data accumulation provided by next-generation sequencing technology has been rapid in recent years. Different varieties of soybean genome were sequenced by several institutes, and these studies will further drive the development of functional markers.

Since within cultivar variation can be quite high to integrate data across experiments, common reference genotypes need to be used in all studies ( Lam et al. 2010 ). Both mutant and QTL association studies are facilitated by the ability to produce inbreds ( reviewed by Lightfoot 2008 ). In species with weak self-incompatibility mechanisms or a low degree of inbreeding depression, like soybean, NILs derived from heterogeneity in RILs populations or their parents can be compared that are fixed for different functional marker alleles. Therefore, the identification of small functional marker effects will not be complicated by polygenic variation. ....

版权说明

本站所提供下载的PDF图书仅提供预览和简介，请支持正版图书。

更多资源请访问:<http://www.tushu007.com>