

<<Barron's AP 物理C>>

图书基本信息

书名：<<Barron's AP 物理C>>

13位ISBN编号：9787510049521

10位ISBN编号：7510049520

出版时间：2012-10

出版公司：世界图书出版公司

作者：(美)佩尔科维茨 等著

页数：671

字数：1115000

版权说明：本站所提供下载的PDF图书仅提供预览和简介，请支持正版图书。

更多资源请访问：<http://www.tushu007.com>

<<Barron's AP 物理C>>

内容概要

备考指南 考点透析

1套诊断测试助您认清强弱项

2套全真测试题帮您考前热身

<<Barron's AP 物理C>>

作者简介

作者：（美国）佩尔科维茨（Robert A. Pelcovits Ph.D.）（美国）法卡斯（Joshua Farkas M.D.）

<<Barron's AP 物理C>>

书籍目录

巴朗五大要点提示

绪论

诊断测试

基础知识

一维运动学

二维运动学

牛顿定律

功、能量和功率

线性动量和质心

转动运动学、力、功和能量

转动惯量、平衡和转动/平衡的结合

简谐运动

万有引力

库仑定律和点电荷产生的电场

由连续电荷分布产生的电场和电势的计算

高斯定律

包含电池的电阻电路分析

.....

章节摘录

版权页：插图：弹性、非弹性和完全非弹性碰撞

1. In an elastic collision, kinetic energy is conserved. Whatever kinetic energy is spent deforming the objects as they collide is completely recovered as they separate (the energy is only temporarily stored as a compression of the objects).

2. In inelastic collisions, kinetic energy is not conserved (some is used to produce heat, sound, or permanently deform the objects, all of which are non-conservative processes).

3. In a totally inelastic collision, kinetic energy is also not conserved, and the two objects remain stuck together after the collision.

碰撞问题的解法 First, you must decide whether the collision is elastic or inelastic. If the problem says that the two particles end up stuck together, it is by definition totally inelastic. Otherwise, the problem generally tells you whether the collision is elastic or (equivalently) whether kinetic energy is conserved. When the collision is elastic, you can construct equations based on the fact that both linear momentum and kinetic energy must be conserved. If the collision is inelastic, you can use only the conservation of linear momentum. The following are extra problem-solving tips:

1. If the collision is totally inelastic, the objects share a common final velocity (because they are stuck together).
2. Remember that momentum is a vector, so if the collision occurs in more than one dimension, the conservation of the momentum vector will be equivalent to a set of two or three one-dimensional equations.
3. If the problem allows you to calculate the change in kinetic energy during the collision, you can use a modified conservation of energy equation (such as $KE = KE_f - KE_0$) to solve an inelastic collision.
4. When you think of a situation where linear momentum is conserved, you probably think of a collision involving "contact forces. However, momentum is conserved even when forces act at a distance (such as gravitational forces or electrostatic forces), as long as there are no net external forces. For example, the momentum of a system of planets, which attract each other gravitationally, is conserved.

<<Barron's AP 物理C>>

编辑推荐

《出国留学书系,SAT、AP备考书系:Barron's AP 物理C(第3版)(英文)》由世界图书出版公司北京公司出版。

<<Barron's AP 物理C>>

版权说明

本站所提供下载的PDF图书仅提供预览和简介, 请支持正版图书。

更多资源请访问:<http://www.tushu007.com>